

AD \_\_\_\_\_

AWARD NUMBER DAMD17-97-1-7222

TITLE: Cell Growth Arrest Mediated by STAT Proteins in Breast Cancer Cells

PRINCIPAL INVESTIGATOR: Dr. Yoshiki Iwamoto

CONTRACTING ORGANIZATION: Yale University School of Medicine  
New Haven, Connecticut 06520-8047

REPORT DATE: July 1999

TYPE OF REPORT: Annual

PREPARED FOR:

U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20000424 202

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE July 1999		3. REPORT TYPE AND DATES COVERED Annual (1 Jul 98 - 30 Jun 99)
4. TITLE AND SUBTITLE Cell Growth Arrest Mediated by STAT Proteins in Breast Cancer Cells				5. FUNDING NUMBERS DAMD17-97-1-7222
6. AUTHOR(S) Yoshiki Iwamoto				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Yale University School of Medicine New Haven, Connecticut 06520-8047  E*Mail: iwamoto@neuro.mssm.edu				8. PERFORMING ORGANIZATION REPORT NUMBER
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research And Materiel Command ATTN: MCMR-RMI-S 504 Scott Street Fort Detrick, Maryland 21702-5012				10. SPONSORING / MONITORING AGENCY REPORT NUMBER
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 words)  We have previously demonstrated that EGF activated STAT1 in some cell lines, causing cell growth arrest and apoptosis. STAT3 is often activated together with STAT1 in response to EGF. It has recently been shown that STAT3 plays an crucial role in the cell cycle progression of BAF/B03 pro-B cells. On the other hand, STAT3 activation by interleukin 6 has been demonstrated to prevent T-cells from apoptosis. More intriguingly, STAT3 but not STAT1 has been reported to be activated constitutively in several breast cancer cell lines and further enhanced in response to EGF. These recent findings led to a new hypothesis that STAT3 activated by EGF may stimulate the cell proliferation competing with activated STAT1, a negative regulator of the cell growth. Therefore, I decided to focus on STAT3 and disrupt its function in mice instead of mutating genome randomly with the frame shift mutagen. To achieve that, I employed the P1 phage-derived Cre/loxP site-specific recombination system since conventional STAT3 knockout mice had been shown to be embryonically lethal. Here I report on generation of mice carrying the loxP-flanked (floxed) (Stat3 <sup>F</sup> ) or -deleted (Stat3 <sup>-</sup> ) mutations in a germline configuration, which has been done.				
14. SUBJECT TERMS Breast Cancer , STAT3, Cre/loxP, Cell Cycle Regulation				15. NUMBER OF PAGES 11
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Y.I. Where copyrighted material is quoted, permission has been obtained to use such material.

Y.I. Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Y.I. Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


☒ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

☒ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

☒ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

☒ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

☐ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

 7/21/99  
PI - Signature Date

## **TABLE OF CONTENTS:**

1. FRONT COVER	1
2. STANDARD FORM 298	2
3. FOREWORD	3
4. TABLE OF CONTENTS	4
5. INTRODUCTION	5
6. BODY	6
7. APPENDICES	11

## **INTRODUCTION:**

We have previously demonstrated that EGF activated STAT1 in some specific cell lines, causing cell growth arrest (1) and apoptosis (2) through gene induction. STAT3 is often activated together with STAT1 in response to EGF (Fig.1). It has recently been shown that STAT3 plays an crucial role in the G1 to S phase transition of BAF/B03 pro-B cells through the downregulation of p21 and p27, and the concomitant upregulation of cyclins D2, D3 and A, and cdc25A (3). On the other hand, STAT3 activation by interleukin 6 (IL-6) has been demonstrated to prevent T-cells from apoptosis (4). More intriguingly, STAT3 but not STAT1 has been reported to be activated constitutively in several breast cancer cell lines and further enhanced in response to EGF (5, 6). We have indeed observed the same phenomenon (Iwamoto, et al., data not shown). These recent findings led me to a new hypothesis that the balance of STAT1 and STAT3 activation could affect the cell growth regulation. Namely, STAT3 activated by EGF may stimulate the cell proliferation competing with activated STAT1, a negative regulator of the cell growth. Based on this hypothesis, I concluded that specific inactivation of STAT3 should be more efficient and straightforward to clarify the molecular mechanism of cell cycle regulation by STAT proteins than inducing random mutations in breast cancer cells as originally proposed in Statement of Work, Task 2. To achieve that, I employed a gene targeting technique with mice. Since conventional STAT3 knockout mice had been shown to be embryonically lethal (7), I decided to use the P1 phage-derived Cre/*loxP* site-specific recombination system (8) and disrupt the Stat3 gene in a mammary gland-specific manner. I have already generated mice carrying the *loxP*-flanked (floxed) (Stat3<sup>F</sup>) or deleted (Stat3<sup>-</sup>) mutations in a germline configuration. To achieve STAT3 inactivation specifically in mammary epithelial cells, I plan to cross Stat3<sup>F</sup> mice with transgenic mice that express Cre under the control of the mouse mammary tumor virus (MMTV) long term repeat (LTR). The MMTV LTR is widely used to express oncogenes such as Ha-ras (9), c-myc (9), c-neu (10), v-src (11) and fgf8 (12) in mouse mammary epithelium to investigate mammary tumorigenesis. I will, then, prepare mammary epithelial primary cultures from both mutant and control mice and investigate the function of STAT proteins in cell cycle regulation and apoptosis as described in my proposal. Here I will report on generation of STAT3-floxed mice which has been done.

## **BODY:**

### *Cloning of genomic Stat3 gene*

Genomic clones containing SH2 domain of the murine Stat3 gene were isolated from a  $\lambda$ Fix II murine 129/Sv genomic library by screening the library using the StuI-ScaI fragment of the murine Stat3 cDNA as a probe. Six phage clones (clones 1, 2, 3, 4, 7 and 10) containing SH2 domain sequences of the murine Stat3 gene were identified.

### *Generation of the Stat3 targeting vector*

The targeting vector was constructed as follows. A 3.8 Kb SmaI fragment and a 2.7 Kb NotI-SmaI fragment of clone 1 (Fig. 2) were inserted into pBluescript KS to generate pBS-1-A and pBS-1-B, respectively. DNA sequence and PCR analyses showed that pBS-1-A contained exons 20 (3' end), 21 and 22, and that pBS-1-B contained exons 15, 16, 17, 18, 19 and 20 (5' end). SH2 domain was found to be coded by exons 19 and 20. We planned to delete SH2 domain specifically by Cre-loxP site-dependent recombination. A loxP site coupled with a HindIII restriction site was introduced into intron 20 of pBS-1-A by insertional mutagenesis using Transformer Site-Directed Mutagenesis Kit (CLONTECH, Palo Alto, CA) (Construct I). Because introns 18 and 19 are too short to introduce another loxP site without regard to interference of splicing, we produced a BglII restriction site in intron 17 of pBS-1-B by site-directed mutagenesis using the same kit as above and inserted a neomycin resistance gene cassette (*neo<sup>r</sup>*) flanked by two loxP sites into the BglII site (Construct II). A SmaI fragment was excised from Construct I and inserted into the SmaI site of Construct II to get Construct III. The pBluescript SK carrying two tandem copies of HSV-thymidine kinase gene cassettes (*HSV-tk*) were digested with NotI and XhoI to recover the fragment which contains *HSV-tks* with the vector. This fragment was replaced with the vector portion of Construct III using the NotI and XhoI sites to get the final construct, pNT-ST3 (Fig. 2).

### *Transfection of embryonic stem cells and generation of mice*

The R1 embryonic stem cell line (13) has been cultured as described. Embryonic stem cells were electroporated with 25  $\mu$ g of NotI-linearized pNT-ST3 per  $1 \times 10^7$  cells and grown under double selection as described (14). We analyzed 112 embryonic stem cell colonies for homologous recombination by Southern blotting. A genomic DNA was digested with HindIII, and two kinds of probes referred to as A and B were used. Nine clones out of 112 were positive for desirable recombination (Fig. 2). Two positive clones (clones 21 and 31) were microinjected into blastocysts of B6 mice to generate mice carrying the heterozygous mutation in the germ line (Stat3<sup>F-Neo/+</sup>).

### *Removal of *neo<sup>r</sup>* from the mutant allele (Stat3<sup>F-Neo</sup>) in vivo*


Removal of *neo<sup>r</sup>* from the mutant allele is desirable to get a higher efficiency of Cre-dependent recombination *in vivo*. *Neo<sup>r</sup>* may also exert cytotoxicity dependently on the genomic locus it is integrated in. To remove *neo<sup>r</sup>* *in vivo*, we first crossed a Stat3<sup>F-Neo/+</sup> mouse with the transgenic mouse expressing Cre recombinase systemically including germ cells (transgenic line TTR 19-5; Iwamoto et al., unpublished) to get the Stat3<sup>F-Neo/+</sup> mouse carrying the transgene. The resulting mouse (Stat3<sup>F-Neo/+</sup>; Cre<sup>+/+</sup>) was crossed with the Stat3<sup>F-Neo/+</sup> mouse. The offsprings were then analyzed for Cre-mediated recombination by PCR as described below. Two offsprings out of eleven contained in the germ line the mutant allele from which only *neo<sup>r</sup>* had been removed (Stat3<sup>F</sup>) (Fig. 3). The rest of all nine offsprings had the mutant allele from which both *neo<sup>r</sup>* and SH2 domain had been deleted (Stat3<sup>-</sup>) (Fig. 3). Alternatively, instead of TTR 19-5, we used another Cre transgenic line, Splicer (Koni et al., unpublished), that expresses the transgene exclusively in ovaries. The same types of Cre-mediated recombination were observed with the similar frequencies.

*Detection of wild-type, Stat3<sup>F-Neo</sup>, Stat3<sup>F</sup> and Stat3<sup>-</sup> mutant alleles by polymerase chain reaction*

Genomic DNA isolated from mouse tails was subjected to PCR for genotyping. The primer pairs a (5'-TATCACTTGCTCCACCTGCTGG-3' and 5'-GCTGGCTCATAGGCAAAAACAC-3'), b (5'-GTTGCTAAGTGTGTTTGAAGAGC-3' and 5'-ACATGTACTTACAGGGTGTGTGC-3'), and c (5'-GTTGCTAAGTGTGTTTGAAGAGC-3' and 5'-GCTGGCTCATAGGCAAAAACAC-3') were used to detect Stat<sup>F</sup>, Stat<sup>F</sup> and wild-type, and Stat3<sup>-</sup> alleles, respectively (Fig.3). The allele carrying the *neo<sup>r</sup>* cassette can be detected by the primer pair d (5'-CCTCAGAAGAACTCGTCAAGAAGG-3' and 5'-GCCAATATGGGATCGGGCCATTGA-3'). The Stat3<sup>F-Neo</sup> allele can be identified as the one which are positive for both Stat<sup>F</sup> and *neo<sup>r</sup>* (Fig.3).

*References:*


1. Chin, Y. E., M. Kitagawa, W. C. Su, Z. H. You, Y. Iwamoto, and X. Y. Fu. 1996. Cell growth arrest and induction of cyclin-dependent kinase inhibitor p21 WAF1/CIP1 mediated by STAT1. *Science* **272**:719-22.
2. Chin, Y. E., M. Kitagawa, K. Kuida, R. A. Flavell, and X. Y. Fu. 1997. Activation of the STAT signaling pathway can cause expression of caspase 1 and apoptosis. *Mol Cell Biol* **17**:5328-37.
3. Fukada, T., T. Ohtani, Y. Yoshida, T. Shirogane, K. Nishida, K. Nakajima, M. Hibi, and T. Hirano. 1998. STAT3 orchestrates contradictory signals in cytokine-induced G1 to S cell-cycle transition. *Embo J* **17**:6670-7.
4. Takeda, K., T. Kaisho, N. Yoshida, J. Takeda, T. Kishimoto, and S. Akira. 1998. Stat3 activation is responsible for IL-6-dependent T cell proliferation through preventing apoptosis: generation and characterization of T cell- specific Stat3-deficient mice. *J Immunol* **161**:4652-60.
5. Garcia, R., C. L. Yu, A. Hudnall, R. Catlett, K. L. Nelson, T. Smithgall, D. J. Fujita, S. P. Ethier, and R. Jove. 1997. Constitutive activation of Stat3 in fibroblasts transformed by diverse oncoproteins and in breast carcinoma cells. *Cell Growth Differ* **8**:1267-76.
6. Watson, C. J. and W. R. Miller. 1995. Elevated levels of members of the STAT family of transcription factors in breast carcinoma nuclear extracts. *Br J Cancer* **71**: 840-4.
7. Takeda, K., K. Noguchi, W. Shi, T. Tanaka, M. Matsumoto, N. Yoshida, T. Kishimoto, and S. Akira. 1997. Targeted disruption of the mouse Stat3 gene leads to early embryonic lethality. *Proc Natl Acad Sci U S A* **94**:3801-4.
8. Gu, H., J. D. Marth, P. C. Orban, H. Mossmann, and K. Rajewsky. 1994. Deletion of a DNA polymerase beta gene segment in T cells using cell type-specific gene targeting [see comments]. *Science* **265**:103-6.
9. Sinn, E., W. Muller, P. Pattengale, I. Tepler, R. Wallace, and P. Leder. 1987. Coexpression of MMTV/v-Ha-ras and MMTV/c-myc genes in transgenic mice: synergistic action of oncogenes in vivo. *Cell* **49**:465-75.
10. Muller, W. J., E. Sinn, P. K. Pattengale, R. Wallace, and P. Leder. 1988. Single-step induction of mammary adenocarcinoma in transgenic mice bearing the activated c-neu oncogene. *Cell* **54**:105-15.
11. Smith, P. D., and M. R. Crompton. 1998. Expression of v-src in mammary epithelial cells induces transcription via STAT3. *Biochem J* **331**:381-5.
12. Daphna-Iken, D., D. B. Shankar, A. Lawshe, D. M. Ornitz, G. M. Shackleford, and C. A. MacArthur. 1998. MMTV-Fgf8 transgenic mice develop mammary and salivary gland neoplasia and ovarian stromal hyperplasia. *Oncogene* **17**:2711-7.
13. Hogan, B., R. Beddington, F. Costantini, and E. Lacy. 1994. *Manipulating the mouse embryo*. Cold Spring Harbor Laboratory Press, New York.
14. Mansour, S.L., K.R. Thomas, and M.R. Capecchi. 1988. Disruption of the proto-oncogene int-2 in mouse embryo-derived stem cells: a general strategy for targeting mutations to non-selectable genes. *Nature* **336**:348-52.




**Figure 1. Activation of STAT proteins by EGF in A431 cells.**

A431 cells were deprived of serum for 17 h and stimulated with 100 ng/ml EGF or 1 ng/ml interferon- $\gamma$  for 15 min. One microliter of the whole cell extract was subjected to EMSA using M-67 SIE as a probe. Three types of the STAT protein-SIE complexes (SIF-A, B and C) were observed. SIF: STAT3 homodimer; SIF-B: STAT1/STAT3 heterodimer; SIF-C: STAT1 homodimer.





**Figure 2. Targeting of loxP sites into the Stat3 locus of ES cells (R1)**



**Figure 3 . PCR genotype analysis of the Stat3 gene.**

**Left panel: Agarose gel electrophoresis of PCR products.**

The genotypes for the samples are Stat3<sup>F/+</sup> (lane 2), Stat3<sup>-/+</sup> (lane 3), Stat3<sup>+/+</sup> (lane 4) and Stat3<sup>F-Neo/+</sup> (lane 5). The letters a, b, c and d beside the panel demonstrate pairs of PCR primers used for each reaction. These primers were designed against the regions represented in the right panel.

**Right panel: Schematic representation of wild-type and mutant Stat3 alleles.**

The PCR product from the wild-type Stat3 allele (WT) is observed in lanes 2-5.

The products from Stat3<sup>F</sup>, Stat3<sup>-</sup> and Stat3<sup>F-Neo</sup> alleles are observed in lanes 2, 3 and 5, respectively.

## **APPENDICES:**

### **1) Key research accomplishments:**

- a. Cloning of genomic Stat3 gene
- b. Making the targeting vector, pNT-ST3
- c. Selection of ES cell clones which are positive for homologous recombination
- d. Generation of the transgenic line (TTR 19-5) expressing Cre recombinase systemically
- e. Generation of Stat3-floxed (Stat3F) and -deleted (Stat3-) mice

### **2) Reportable outcomes:**

#### **i) Developed reagents:**

- a. ES cell lines were developed that carry a floxed-mutation in the Stat3 genomic locus
- b. The transgenic line (TTR 19-5) expressing Cre recombinase systemically
- c. Stat3-floxed (Stat3F) and -deleted (Stat3-) mice

#### **ii) Funding applied for based on work supported by this award:**

- a. Career Development Awards of U.S. Army 1999 Breast Cancer Research Program

#### **iii) Employment received based on experiences/training supported by this award:**

- a. I was offered a position of Research Assistant Professor at the Mount Sinai School of Medicine.